L18 ANSWER 2 OF 8 MEDLINE DUPLICATE 1

ACCESSION NUMBER: 2001610635 MEDLINE

DOCUMENT NUMBER: 21541943 PubMed ID: 11590405

TITLE: Dendritic cell maturation is required for the

cross-tolerization of CD8+ T cells.

COMMENT: Comment in: Nat Immunol. 2001 Nov;2(11):988-9

AUTHOR: Albert M L; Jegathesan M; Darnell R B

CORPORATE SOURCE: Laboratory of Neuro-Oncology, The Rockefeller University,

1230 York Avenue, Box 26, New York, NY 10021, USA..

albertm@rockvax.rockefeller.edu

SOURCE: Nat Immunol, (2001 Nov) 2 (11) 1010-7.

Journal code: 100941354. ISSN: 1529-2908.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 200112

ENTRY DATE: Entered STN: 20011102

Last Updated on STN: 20020123

Entered Medline: 20011213

AB In vivo models have shown that tissue-restricted antigen may be captured by bone marrow-derived cells and cross-presented for the

tolerization of CD8+ T cells. Although these studies have shown peripheral tolerization of CD8+ T cells, the mechanism of antigen transfer and the nature of the antigen-presenting cell (APC) remain undefined. We report here the establishment of an in vitro system for the study of cross-tolerance and show that dendritic cells (DCs) phagocytose apoptotic cells and tolerize antigen-specific CD8+ T cells when cognate CD4+ T helper cells are absent. Using this system, we directly tested the "two-signal" hypothesis for the regulation of priming versus tolerance. We found that the same CD83+ myeloid-derived DCs were required for both cross-priming and cross-tolerance. These data suggested that the current model for peripheral T cell tolerance, "signal 1 in the absence of signal 2", requires refinement: the critical checkpoint is not DC maturation, but instead the presence of a third signal, which is

active

at the DC-CD4+ T cell interface.

18 aptamer

142, 25, 9

1<sup>1</sup>/<sub>1</sub><sup>2</sup> 9 6 4 5

lym

, v, 1